

Product datasheet for **TA387660**

H2BC4 Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IF, IP, WB
Recommended Dilution:	Recommended dilution: WB:1:500-1:2000, IF:1:1-1:10, IP:1:200-1:2000
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Peptide sequence around site of Acetyl-Lys (15) derived from Human Histone H2B type 1-C/E/F/G/I
Formulation:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Concentration:	lot specific
Purification:	Antigen Affinity Purified
Conjugation:	Unconjugated
Storage:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Stability:	1 year from dispatch.
Database Link:	P62807
Background:	Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.




[View online »](#)

Product images:



Immunofluorescence staining of HeLa cells (treated with 30mM sodium butyrate for 4h) with TA387660 at 1:5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).




Immunoprecipitating HIST1H2BC in HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h)

Lane 1: Rabbit control IgG instead of TA387660 in HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h). For western blotting, a HRP-conjugated Protein G antibody was used as the secondary antibody (1/2000)

Lane 2: TA387660 (5µg) + HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h) (500µg)

Lane 3: HepG2 whole cell lysate (treated with 30mM sodium butyrate for 4h) (20µg)



Western Blot

Detected samples: HeLa whole cell lysate, K562 whole cell lysate, 293 whole cell lysate; Untreated (-) or treated (+) with 30mM sodium butyrate for 4h

All lanes: HIST1H2BC antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1/40000 dilution

Predicted band size: 14 kDa

Observed band size: 14 kDa