

Product datasheet for TA387859M

MGAT4B Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Alpha-1,3-mannosyl-glycoprotein 4-beta-N-

acetylglucosaminyltransferase B protein (380-522AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Concentration: lot specific

Purification: >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: Q9UQ53

Background: Glycosyltransferase that participates in the transfer of N-acetylglucosamine (GlcNAc) to the

core mannose residues of N-linked glycans. Catalyzes the formation of the GlcNAcbeta1-4 branch on the GlcNAcbeta1-2Manalpha1-3 arm of the core structure of N-linked glycans. Essential for the production of tri- and tetra-antennary N-linked sugar chains. Has lower affinities for donors or acceptors than MGAT4A, suggesting that, under physiological

conditions, it is not the main contributor in N-glycan biosynthesis.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Western Blot
Positive WB detected in: SH-SY5Y whole cell
lysate, Jurkat whole cell lysate
All lanes: MGAT4B antibody at 3µg/ml
Secondary
Goat polyclonal to rabbit IgG at 1/50000 dilution
Predicted band size: 64, 65 kDa
Observed band size: 64 kDa



IHC image of [TA387859] diluted at 1:300 and staining in paraffin-embedded human kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.