

Product datasheet for **TA387948**


CYP4F11 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	IF, WB
Recommended Dilution:	Recommended dilution: WB:1:1000-1:5000, IF:1:50-1:200
Reactivity:	Mouse, Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant Human Phylloquinone omega-hydroxylase CYP4F11 protein (38-524AA)
Formulation:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Concentration:	lot specific
Purification:	>95%, Protein G purified
Conjugation:	Unconjugated
Storage:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Stability:	1 year from dispatch.
Database Link:	Q9HBI6
Background:	Omega-hydroxylase that oxidizes a variety of structurally unrelated compounds, including fatty acids and xenobiotics. Plays a key role in vitamin K catabolism by mediating omega-hydroxylation of vitamin K1 (phylloquinone), and menaquinone-4 (MK-4), a form of vitamin K2. Hydroxylation of phylloquinone and MK-4 probably regulates blood coagulation (PubMed:24138531). Catalyzes omega-hydroxylation of 3-hydroxy fatty acids, such as 3-hydroxypalmitate, 3-hydroxyoleate, 3-hydroxyarachidonate, and 3-hydroxystearate (PubMed:18065749, PubMed:19932081). Oxidizes drugs such as erythromycin, benzphetamine, ethylmorphine, chlorpromazine and imipramine.



[View online »](#)

Product images:

Immunofluorescent analysis of HepG2 cells using TA387948 at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L)



Western blot
All lanes: CYP4F11 antibody at 2 μ g/ml
Lane 1: Mouse liver tissue
Lane 2: Mouse kidney tissue
Secondary
Goat polyclonal to rabbit IgG at 1/10000 dilution
Predicted band size: 61 kDa
Observed band size: 61 kDa