

Product datasheet for **TA387997M**

ATP6V1G2 Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IF, IHC, WB
Recommended Dilution:	Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200, IF:1:50-1:200
Reactivity:	Mouse, Rat, Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant Human V-type proton ATPase subunit G 2 protein (1-118AA)
Formulation:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Concentration:	lot specific
Purification:	>95%, Protein G purified
Conjugation:	Unconjugated
Storage:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Stability:	1 year from dispatch.
Database Link:	Q95670
Background:	Catalytic subunit of the peripheral V1 complex of vacuolar ATPase (V-ATPase). V-ATPase is responsible for acidifying a variety of intracellular compartments in eukaryotic cells.




[View online »](#)


Product images:



Immunohistochemistry of paraffin-embedded human brain tissue using [TA387997] at dilution of 1:100



Immunofluorescence staining of A549 cells with [TA387997] at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Western blot

All lanes: ATP6V1G2 antibody at 12µg/ml

Lane 1: Rat heart tissue

Lane 2: Mouse spleen tissue

Lane 3: Hela whole cell lysate

Lane 4: HepG2 whole cell lysate

Lane 5: A549 whole cell lysate

Lane 6: HT29 whole cell lysate

Lane 7: K562 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 14, 9 kDa

Observed band size: 14 kDa