

Product datasheet for TA388019M

Product data:

Product Type: Primary Antibodies

H1-4 Rabbit Polyclonal Antibody

Applications: IF, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:2000, IF:1:1-1:10

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide sequence around site of Phospho-Thr (17) derived from Human Histone H1.4

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Concentration: lot specific

Purification: Antigen Affinity Purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: P10412

Background: Histone H1 protein binds to linker DNA between nucleosomes forming the macromolecular

structure known as the chromatin fiber. Histones H1 are necessary for the condensation of nucleosome chains into higher-order structured fibers. Acts also as a regulator of individual gene transcription through chromatin remodeling, nucleosome spacing and DNA methylation

(By similarity).




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:



Immunofluorescence staining of Hela cells with [TA388019] at 1:2.5, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).



Western Blot

Detected samples: Jurkat whole cell lysate, Hela whole cell lysate; Untreated (-) or treated (+) with

Calyculin A and EGF

All lanes: HIST1H1E antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1/40000 dilution

Predicted band size: 22 kDa Observed band size: 22 kDa