

Product datasheet for TA388146

THNSL2 Rabbit Polyclonal Antibody

Product data:

OriGene Technologies, Inc.


9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product Type:Primary AntibodiesApplications:IHC, WBRecommended Dilution:Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500Reactivity:HumanHost:RabbitIsotype:IgGClonality:PolyclonalImmunogen:Recombinant Human Threonine synthase-like 2 protein (295-441AA)Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Forneutation:Id specificPurification:José Protein G purifiedForngues:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:Jear from dispatch.Database Link:Osóryf1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and phosphate (By similarity).		
Recommended Dilution:Recommended dilution:WB:1:500-1:5000, IHC:1:200-1:500Reactivity:HumanHost:RabbitIsotype:IgGIsotype:lgGClonality:PolyclonalImmunogen:Recombinant Human Threonine synthase-like 2 protein (295-441AA)Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:Iot specificPurification:Sp5%, Protein G purifiedConjugation:UnconjugatedStability:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:Q86YI6Background:Soform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Product Type:	Primary Antibodies
Reactivity:HumanHost:RabbitIsotype:IgGClonality:PolyclonalImmunogen:Recombinant Human Threonine synthase-like 2 protein (295-441AA)Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:Iot specificPurification:Jostph Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YJ6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Applications:	IHC, WB
Host:RabbitHost:RabbitIsotype:IgGClonality:PolyclonalImmunogen:Recombinant Human Threonine synthase-like 2 protein (295-441AA)Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:lot specificPurification:>95%, Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YI6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Recommended Dilution:	Recommended dilution: WB:1:500-1:5000, IHC:1:200-1:500
Isotype:IgGIsotype:lgGClonality:PolyclonalImmunogen:Recombinant Human Threonine synthase-like 2 protein (295-441AA)Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:lot specificPurification:>95%, Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YJ6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Reactivity:	Human
Clonality:PolyclonalImmunogen:Recombinant Human Threonine synthase-like 2 protein (295-441AA)Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:lot specificPurification:>95%, Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YJ6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Host:	Rabbit
Immunogen:Recombinant Human Threonine synthase-like 2 protein (295-441AA)Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:lot specificPurification:>95%, Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YI6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	lsotype:	IgG
Formulation:Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:lot specificPurification:>95%, Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YI6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Clonality:	Polyclonal
Constituents: 50% Glycerol, 0.01M PBS, pH 7.4Concentration:lot specificPurification:>95%, Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YI6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Immunogen:	Recombinant Human Threonine synthase-like 2 protein (295-441AA)
Purification:>95%, Protein G purifiedConjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YI6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Formulation:	
Conjugation:UnconjugatedStorage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YI6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Concentration:	lot specific
Storage:Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.Stability:1 year from dispatch.Database Link:Q86YI6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Purification:	>95%, Protein G purified
Stability:1 year from dispatch.Database Link:Q86YJ6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Conjugation:	Unconjugated
Database Link:Q86YJ6Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Storage:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Background:Isoform 1: Acts as a catabolic phospho-lyase on both gamma- and beta-phosphorylated substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Stability:	1 year from dispatch.
substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and	Database Link:	<u>Q86YJ6</u>
	Background:	substrates. Degrades O-phospho-threonine (PThr) to alpha-ketobutyrate, ammonia and




This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US

Product images:



IHC image of TA388146 diluted at 1:400 and staining in paraffin-embedded human ovarian cancer performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Western Blot Positive WB detected in: Hela whole cell lysate, HepG2 whole cell lysate All lanes: THNSL2 antibody at 4.3µg/ml Secondary Goat polyclonal to rabbit lgG at 1/50000 dilution Predicted band size: 55, 46, 43, 48, 28 kDa Observed band size: 55 kDa

This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US