

Product datasheet for **TA388292M**

TRMT6 Rabbit Polyclonal Antibody


Product data:

| | |
|-----------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | IHC, WB |
| Recommended Dilution: | Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200 |
| Reactivity: | Human |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Recombinant Human tRNA (adenine(58)-N(1))-methyltransferase non-catalytic subunit TRM6 protein (1-300AA) |
| Formulation: | Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4 |
| Concentration: | lot specific |
| Purification: | >95%, Protein G purified |
| Conjugation: | Unconjugated |
| Storage: | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze. |
| Stability: | 1 year from dispatch. |
| Database Link: | Q9UJA5 |
| Background: | Substrate-binding subunit of tRNA (adenine-N(1))-methyltransferase, which catalyzes the formation of N(1)-methyladenine at position 58 (m1A58) in initiator methionyl-tRNA. |




[View online »](#)


Product images:



Immunohistochemistry of paraffin-embedded human testis tissue using [TA388292] at dilution of 1:100



Immunohistochemistry of paraffin-embedded human adrenal gland tissue using [TA388292] at dilution of 1:100



Western blot

All lanes: TRMT6 antibody at 4 μ g/ml

Lane 1: U251 whole cell lysate

Lane 2: HepG2 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 56, 45, 41, 37 kDa

Observed band size: 56 kDa