

## **Product datasheet for TA388331M**

## **FBXO6** Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human F-box only protein 6 protein (104-293AA)

**Formulation:** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:** lot specific

**Purification:** Antigen Affinity Purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q9NRD1

**Background:** Substrate-recognition component of some SCF (SKP1-CUL1-F-box protein)-type E3 ubiquitin

ligase complexes. Involved in endoplasmic reticulum-associated degradation pathway (ERAD)

for misfolded lumenal proteins by recognizing and binding sugar chains on unfolded glycoproteins that are retrotranlocated into the cytosol and promoting their ubiquitination and subsequent degradation. Able to recognize and bind denatured glycoproteins, which are

modified with not only high-mannose but also complex-type oligosaccharides. Also recognizes sulfated glycans. Also involved in DNA damage response by specifically

recognizing activated CHEK1 (phosphorylated on 'Ser-345'), promoting its ubiquitination and degradation. Ubiquitination of CHEK1 is required to insure that activated CHEK1 does not accumulate as cells progress through S phase, or when replication forks encounter transient

impediments during normal DNA replication.




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Immunohistochemistry of paraffin-embedded human glioma using [TA388331] at dilution of 1:100



Immunohistochemistry of paraffin-embedded human kidney tissue using [TA388331] at dilution of 1:100





Western blot

All lanes: FBXO6 antibody at 3.87 µg/ml + A431

whole cell lysate Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 34 kDa Observed band size: 34, 120 kDa