

Product datasheet for TA388333M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

MTMR2 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:300-1:500, IF:1:50-1:200

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human Myotubularin-related protein 2 protein (1-173AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, pH 7.4

Concentration: lot specific

Purification: >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.


Database Link: Q13614

Background: Phosphatase that acts on lipids with a phosphoinositol headgroup. Has phosphatase activity


towards phosphatidylinositol 3-phosphate and phosphatidylinositol 3,5-bisphosphate.



Product images:




IHC image of [TA388333] diluted at 1:400 and staining in paraffin-embedded human testis tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a biotinylated secondary antibody and visualized using an HRP conjugated SP system.



Immunofluorescence staining of Hela cells with [TA388333] at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-congugated AffiniPure Goat Anti-Rabbit IgG(H+L).





Western Blot

Positive WB detected in: Hela whole cell lysate All lanes: MTMR2 antibody at 5.5µg/ml Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 74, 66 kDa Observed band size: 74 kDa