

## **Product datasheet for TA388413**

## **PPIAL4G Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** IHC, WB

Recommended Dilution: Recommended dilution: WB:1:500-1:5000, IHC:1:20-1:200

Reactivity: Mouse, Human

Rabbit Host: Isotype: lgG

Polyclonal Clonality:

Immunogen: Recombinant Human Peptidyl-prolyl cis-trans isomerase A-like 4G protein (1-164AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Concentration: lot specific

**Purification:** >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: P0DN37

Background: PPlases accelerate the folding of proteins. It catalyzes the cis-trans isomerization of proline

imidic peptide bonds in oligopeptides.




OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Western Blot


All lanes: PPIAL4G antibody at 6µg/ml

Lane 1: Mouse liver tissue Lane 2: Mouse spleen tissue Lane 3: Mouse lung tissue Lane 4: Mouse kidney tissue

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 19 kDa Observed band size: 19 kDa



Immunohistochemistry of paraffin-embedded human kidney tissue using TA388413 at dilution of 1:100