

Product datasheet for TA388542

OR6B3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:100-1:500

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Peptide sequence from Human Olfactory receptor 6B3 protein (259-272AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

Concentration: lot specific

Purification: >95%, Protein G purified

Conjugation: Unconjugated

Storage: Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

Stability: 1 year from dispatch.

Database Link: Q8NGW1

Background: Odorant receptor.


OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Product images:




IHC image of TA388542 diluted at 1:300 and staining in paraffin-embedded human skin tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



IHC image of TA388542 diluted at 1:300 and staining in paraffin-embedded human brain tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.





Western Blot

Positive WB detected in: U251 whole cell lysate All lanes: OR6B3 antibody at 1:2000 Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 38 kDa Observed band size: 38 kDa