

Product datasheet for **TA388570M**

RPUSD4 Rabbit Polyclonal Antibody


Product data:

Product Type:	Primary Antibodies
Applications:	IHC, WB
Recommended Dilution:	Recommended dilution: WB:1:500-1:2000, IHC:1:20-1:200
Reactivity:	Human
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Recombinant Human Mitochondrial RNA pseudouridine synthase RPUSD4 protein (41-240AA)
Formulation:	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, PH 7.4
Concentration:	lot specific
Purification:	Antigen Affinity Purified
Conjugation:	Unconjugated
Storage:	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Stability:	1 year from dispatch.
Database Link:	Q96CM3
Background:	Catalyzes uridine to pseudouridine isomerization (pseudouridylation) of different mitochondrial RNA substrates. Acts on position 1397 in 16S mitochondrial ribosomal RNA (16S mt-rRNA). This modification is required for the assembly of 16S mt-rRNA into a functional mitochondrial ribosome (PubMed:27974379). Acts on position 39 in mitochondrial tRNA(Phe) (PubMed:28082677). As a component of a functional protein-RNA module, consisting of RCC1L, NGRN, RPUSD3, RPUSD4, TRUB2, FASTKD2 and 16S mt-rRNA, controls 16S mt-rRNA abundance and is required for intra-mitochondrial translation (PubMed:27667664).




[View online »](#)

Product images:



Immunohistochemistry of paraffin-embedded human colon cancer using [TA388570] at dilution of 1:100



Immunohistochemistry of paraffin-embedded human kidney tissue using [TA388570] at dilution of 1:100



Western blot

All lanes: RPUSD4 antibody at 0.5µg/ml + MCF-7 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 43, 39 kDa

Observed band size: 43 kDa