

## **Product datasheet for TA388594M**

## **COPS4 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: Recommended dilution: WB:1:200-1:1000, IHC:1:20-1:200

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant Human COP9 signalosome complex subunit 4 protein (1-310AA)

**Formulation:** PBS with 0.02% sodium azide, 50% glycerol, pH7.3.

**Concentration:** lot specific

**Purification:** Antigen Affinity Purified

Conjugation: Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q9BT78

**Background:** Component of the COP9 signalosome complex (CSN), a complex involved in various cellular

and developmental processes. The CSN complex is an essential regulator of the ubiquitin (UbI) conjugation pathway by mediating the deneddylation of the cullin subunits of SCF-type E3 ligase complexes, leading to decrease the UbI ligase activity of SCF-type complexes such as SCF, CSA or DDB2. Also involved in the deneddylation of non-cullin subunits such as STON2.

The complex is also involved in phosphorylation of p53/TP53, c-jun/JUN,

IkappaBalpha/NFKBIA, ITPK1, IRF8/ICSBP and SNAPIN, possibly via its association with CK2 and PKD kinases. CSN-dependent phosphorylation of TP53 and JUN promotes and protects

degradation by the Ubl system, respectively.




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**




Immunohistochemistry of paraffin-embedded human breast cancer using [TA388594] at dilution of 1:100



Immunohistochemistry of paraffin-embedded human pancreatic tissue using [TA388594] at dilution of 1:100





Western blot

All lanes: COPS4 antibody at 8µg/ml Lane 1: Hela whole cell lysate Lane 2: HepG2 whole cell lysate

Secondary

Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 47, 41 kDa Observed band size: 47 kDa