

## **Product datasheet for TA388662**

## CYB5R2 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:20-1:200

**Reactivity:** Mouse, Human

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human NADH-cytochrome b5 reductase 2 protein (1-237AA)

**Formulation:** Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

**Concentration:** lot specific

**Purification:** >95%, Protein G purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: Q6BCY4

Background: NADH-cytochrome b5 reductases are involved in desaturation and elongation of fatty acids,

cholesterol biosynthesis, drug metabolism, and, in erythrocyte, methemoglobin reduction. Responsible for NADH-dependent lucigenin chemiluminescence in spermatozoa by reducing both lucigenin and 2-[4-iodophenyl]-3-[4-nitrophenyl]-5-[2,4-disulfophenyl]-2H tetrazolium

monosodium salt (WST-1).




**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



Immunohistochemistry of paraffin-embedded human testis tissue using TA388662 at dilution of 1:100



Western blot All lanes: CYB5R2 antibody at 4µg/ml + Mouse kidney tissue Secondary Goat polyclonal to rabbit IgG at 1/10000 dilution

Predicted band size: 32, 28 kDa Observed band size: 32 kDa