

## **Product datasheet for TA388707**

## **Ren1 Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: IHC, WB

Recommended Dilution: Recommended dilution: WB:1:1000-1:5000, IHC:1:100-1:500

Reactivity: Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant Rat Ren1 protein (65-264AA)

Formulation: Preservative: 0.03% Proclin 300

Constituents: 50% Glycerol, 0.01M PBS, PH 7.4

**Concentration:** lot specific

**Purification:** >95%, Antigen Affinity purified & Affinity purified

**Conjugation:** Unconjugated

**Storage:** Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.

**Stability:** 1 year from dispatch.

Database Link: P08424

**Background:** Renin is a highly specific endopeptidase, whose only known function is to generate

angiotensin I from angiotensinogen in the plasma, initiating a cascade of reactions that produce an elevation of blood pressure and increased sodium retention by the kidney.


**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



## **Product images:**



IHC image of TA388707 diluted at 1:100 and staining in paraffin-embedded rat kidney tissue performed on a Leica BondTM system. After dewaxing and hydration, antigen retrieval was mediated by high pressure in a citrate buffer (pH 6.0). Section was blocked with 10% normal goat serum 30min at RT. Then primary antibody (1% BSA) was incubated at 4°C overnight. The primary is detected by a Goat anti-rabbit polymer IgG labeled by HRP and visualized using 0.05% DAB.



Western Blot

Positive WB detected in: Rat Heart tissue, Rat

Liver tissue

All lanes: Ren1 antibody at 1:1000

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 45 kDa Observed band size: 45 kDa