

Product datasheet for **TA388902**

IL12A , IL12B Mouse Monoclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Sandwich ELISA: In a sandwich ELISA (assuming 100ul/well), a concentration of 2.0-4.0 µg/ml of this antibody will detect at least 1000pg/ml Recombinant Human IL-12 when used with ProSci's Biotinylated Antigen Affinity Purified Anti-Human IL-12 as the detection antibody at a concentration of at least 0.25-0.5 µg/ml. Western Blot To detect Human IL-12 by Western Blot analysis this antibody can be used at a concentration of 0.50-1.0 µg/ml. When used in conjunction with compatible secondary reagents the detection limit for Recombinant Human IL-12 is 0.50-1.0 ng/lane, under non-reducing conditions.
Reactivity:	Human
Host:	Mouse
Clonality:	Monoclonal
Immunogen:	Produced in BALB/c x ICR F1 mice immunized with highly pure Recombinant Human IL-12. Anti-Human IL- 12-specific antibody was purified from ascites fluid by Protein A affinity chromatography.
Concentration:	lot specific
Purification:	IL- 12-specific antibody was purified from ascites fluid by Protein A affinity chromatography
Conjugation:	Unconjugated
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Background:	IL-12 is a potent regulator of cell-mediated immune responses and it induces IFN-gamma production by NK and T cells. It is produced by activated monocytes/macrophage cells, B lymphocytes and connective tissue-type mast cells. Among its biological activities, IL-12 promotes the growth and activity of activated NK, CD4+ and CD8+ cells, and induces the development of IFN-gamma-producing Th1 cells. Recombinant Human IL-12 is a 75.0 kDa heterodimeric glycoprotein consisting of disulfide-linked 35 kDa (p35) and 40 kDa (p40) subunits (503 total amino acid residues).



[View online »](#)