

Product datasheet for TA388920

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

II3 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: ELISA, WB

Recommended Dilution: Sandwich ELISA: To detect mIL-3 by sandwich ELISA (using 100 ul/well antibody solution) a

concentration of 0.25 – 1.0 µg/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with ProSci's Polyclonal Anti-Murine IL-3 as a capture antibody,

allows the detection of at least 0.2 – 0.4 ng/well of recombinant mIL-3.

Western Blot

To detect mIL-3 by Western Blot analysis this antibody can be used at a concentration of 0.1 - 0.2 ug/ml. Used in conjunction with compatible secondary reagents the detection limit for recombinant mIL-3 is 1.5 - 3.0 ng/lane, under either reducing or non-reducing conditions.

Reactivity: Mouse
Host: Rabbit
Clonality: Polyclonal

Immunogen: Produced from sera of rabbits immunized with highly pure Recombinant Murine IL-3. Anti-

Murine IL-3-specific antibody was purified by affinity chromatography and then biotinylated.

Concentration: lot specific

Purification: IL-3-specific antibody was purified by affinity chromatography and then biotinylated

Conjugation: Biotin

Storage: Store at -20°C as received.

Stability: Stable for 12 months from date of receipt.

Database Link: P01586

Background: IL-3 is a hematopoietic growth factor that promotes the survival, differentiation and

proliferation of committed progenitor cells of the megakaryocyte, granulocyte-macrophage, erythroid, eosinophil, basophil and mast cell lineages. Produced by T cells, mast cells and eosinophils, IL-3 enhances thrombopoiesis, phagocytosis, and antibody-mediated cellular cytotoxicity. Its ability to activate monocytes suggests that IL-3 may have additional

immunoregulatory roles. Many of the IL-3 activities depend upon co-stimulation with other cytokines. IL-3 is a species-specific, variably glycosylated cytokine. Recombinant Murine IL-3 is

a 15.1 kDa globular protein containing 135 amino acid residues.

