

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Product datasheet for TA388950

Cxcl12 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	Sandwich ELISA: To detect Rat SDF-1 alpha by sandwich ELISA (using 100 ul/well), a concentration of $0.25 - 1.0 \mu$ g/ml of this antibody is required. This biotinylated polyclonal antibody, in conjunction with ProSci's Polyclonal Anti-Rat SDF-1 alpha as a capture antibody, allows the detection of at least $0.2 - 0.4$ ng/well of recombinant Rat SDF-1 alpha. Western Blot To detect Rat SDF-1 alpha by Western Blot analysis, this antibody can be used at a concentration of $0.1 - 0.2 \mu$ g/ml. When used in conjunction with compatible development reagents, the detection limit for recombinant Rat SDF-1 alpha is $1.5 - 3.0$ ng/lane, under either reducing or non-reducing conditions.
Reactivity:	Rat
Host:	Rabbit
Clonality:	Polyclonal
Immunogen:	Produced from sera of rabbits immunized with highly pure recombinant Rat SDF-1 alpha. Anti-Rat SDF-1 alpha-specific antibody was purified by affinity chromatography and then biotinylated.
Concentration:	lot specific
Purification:	SDF-1 alpha-specific antibody was purified by affinity chromatography and then biotinylated
Conjugation:	Biotin
Storage:	Store at -20°C as received.
Stability:	Stable for 12 months from date of receipt.
Background:	SDF-1 alpha andbeta are stromal-derived, CXC chemokines that signal through the CXCR4 receptor. SDF-1alpha andbeta chemoattract B and T cells, and have been shown to induce migration of CD34+ stem cells. Additionally, the SDF-1 proteins exert HIV-suppressive activity in cells expressing the CXCR4 receptor. Human and murine SDF-1 proteins act across species. SDF-1 alpha andbeta contain the four highly conserved cysteine residues present in CXC chemokines. The mature SDF-1alpha protein is the result of alternative splicing of the SDF-1 gene and contains 68 amino acid residues. Recombinant Rat SDF-1 alpha is a 7.9 kDa protein containing 68 amino acid residues.



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US