

## **Product datasheet for TA389004**

# OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

### Phospho-Dennd3 (pThr450) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:1000

**WB Brain**: 1:1000

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic phospho-peptide corresponding to amino acid residues surrounding Thr450 of

human DENND3, conjugated to keyhole limpet hemocyanin (KLH).

Specificity: Specific for endogenous levels of the ~142 kDa DENND3 phosphorylated at Thr450. The

immunolabeling is completely eliminated by treatment with  $\lambda$ -phosphatase.

**Formulation:** 10 mM HEPES (pH 7.5), 150 mM NaCl, 100 μg per ml BSA and 50% glycerol.

**Concentration:** lot specific

**Purification:** Antigen Affinity Purified from Pooled Serum

**Conjugation:** Unconjugated

Storage: Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to

presence of 50% glycerol. Stable for at least 1 year at -20°C.

**Stability:** After date of receipt, stable for at least 1 year at -20°C.

Predicted Protein Size: 142

**Gene Name:** DENN domain containing 3

Database Link: A2RUS2

Background: The DENN (differentially expressed in normal and neoplastic cells) domain (DENND) is a

poorly characterized protein module conserved throughout evolution (McPherson et al., 2011). Proteins bearing a DENN domain have recently emerged as the largest family of Rab GEFs. Among these DENN domain proteins, DENND3 is a GEF for Rab12; promoting the exchange of GDP to GTP, converting inactive GDP-bound RAB12 into its active GTP-bound

form (McPherson and Xu, 2015; Matsui et al., 2011).





#### Phospho-Dennd3 (pThr450) Rabbit Polyclonal Antibody - TA389004

Synonyms: KIAA0870

**Note:** Prepared from pooled rabbit serum by affinity purification via sequential chromatography on

phospho and non-phospho peptide affinity columns.