

Product datasheet for TA389047

ALDH1A1 Mouse Antibody [Clone ID: M562]

Product data:

Product Type: Primary Antibodies

Clone Name: M562 Applications: ICC, WB

Recommended Dilution: WB: 1:1000

ICC: 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Immunogen: Clone M562 was generated from a recombinant protein corresponding to amino acids in the

N-terminal region from human ALDH1A1.

Specificity: This antibody antibody detects a 55 kDa* protein on SDS-PAGE immunoblots of human A431,

HepG2, and mouse liver, as well as human recombinant ALDH1A1. The antibody does not

detect ALDH1A3 in PC3 cells, ALDH1A2 in Jurkat cells, and bovine tubulin (55 kDa).

Formulation: PBS + 1 mg/ml BSA, 0.05% NaN3 and 50% glycerol

Concentration: lot specific

Purification: Protein G Purified

Conjugation: Unconjugated

Storage: Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to

presence of 50% glycerol. Stable for at least 1 year at -20°C.

Stability: After date of receipt, stable for at least 1 year at -20°C.

Predicted Protein Size: 55

Database Link: P00352



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



ALDH1A1 Mouse Antibody [Clone ID: M562] - TA389047

Background:

Aldehyde dehydrogenase (ALDH) superfamily is a ubiquitous group of enzymes found in all taxonomic domains. ALDH detoxifies endogenous and exogenous aldehydes, protecting cellular homeostasis and organismal functions. These enzymes are necessary for the synthesis of retinoic acid, betaine, and folate. Recent studies have reported high levels of ALDH found in cancer cells, suggesting that ALDH can act as a marker for cancer cells found in a wide variety of tissues including skin, prostate, lung, and neural tissues. Additionally, certain diseases can be identified when ALDH activity is absent. ALDH1A1 is vital for retinol synthesis and alcohol metabolism. ALDH1A1 active sites include an active cysteine residue, which catalyses the transformation of aldehydes into their respective carboxylic groups. ALDH1A1 amino acid sequence and function is highly conserved in humans and rodents.

Note:

Protein G purified tissue culture supernatant.