

Product datasheet for TA389092

BSG Mouse Antibody [Clone ID: M036]

Product data:

Product Type: Primary Antibodies

Clone Name: M036

Applications: ICC, IP, WB Recommended Dilution: **WB**: 1:2000

ICC: 1:100

Reactivity: Human
Host: Mouse
Isotype: IgG2a

Immunogen: Clone (M036) was generated from a proprietary antigen related to the native human CD147

expressed in NCI-H1915 lung cancer cell line.

Specificity: Clone M036 mouse monoclonal antibody detects a 38-58 kDa* protein on SDS-PAGE

immunoblots of human BEAS, NCIH28, NCI-H1299, LNCaP, and A431 cells, as well as human brain tissues. The antibody works for western blot, immunoprecipitation, ELISA capture, and

immunocytochemistry.

Formulation: PBS + 1 mg/ml BSA, 0.05% NaN3 and 50% glycerol

Concentration: lot specific

Purification: Protein G Purified

Conjugation: Unconjugated

Storage: Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to

presence of 50% glycerol. Stable for at least 1 year at -20°C.

Stability: After date of receipt, stable for at least 1 year at -20°C.

Predicted Protein Size: 38-58

Database Link: P35613



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Background:

D147 (Basigin/EMMPRIN) is a type I integral membrane receptor protein belonging to the immunoglobulin superfamily. CD147 is a glycosylated protein with four known isoforms with isoform 2 being the most abundantly expressed. In many cancers, CD147 is overexpressed and stimulates the secretion of extracellular matrix metalloproteinases by tumor fibroblasts, which can promote tumor progression. CD147 is co-expressed with the amino acid transporter, CD98h, in metabolically active cells such as lymphocytes, macrophages, and tumor cells. CD147 interaction with caveolin and cyclophilins regulates its functions, and this protein may function outside of tumor cells through cell shedding and microvesicle release.

Note:

Protein G purified tissue culture supernatant.