

Product datasheet for **TA389153**

ITGB1 Mouse Antibody [Clone ID: M032]

Product data:

Product Type:	Primary Antibodies
Clone Name:	M032
Applications:	ICC, WB
Recommended Dilution:	WB: 1:500 ICC: 1:200
Reactivity:	Human
Host:	Mouse
Isotype:	IgG1
Immunogen:	Clone M032 was generated from a proprietary antigen related to the extracellular region of human integrin β 1 expressed in A431 epidermoid carcinoma cell line.
Specificity:	Clone M032 detects a 120-150 kDa* band corresponding to Integrin β 1 on SDS-PAGE immunoblots of "Native" human A431, A549, MDA-MB-231, and LNCaP cell lysates, as well as human breast, skin, and lung tissues. The antibody weakly detects the denatured form of Integrin β 1. Clone M032 can be used in western blot, immunocytochemistry, and ELISA.
Formulation:	PBS + 1 mg/ml BSA, 0.05% NaN ₃ and 50% glycerol
Concentration:	lot specific
Purification:	Protein G Purified
Conjugation:	Unconjugated
Storage:	Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to presence of 50% glycerol. Stable for at least 1 year at -20°C.
Stability:	After date of receipt, stable for at least 1 year at -20°C.
Predicted Protein Size:	120-150
Database Link:	P05556



[View online »](#)

Background:

Integrins are cell adhesion molecules that can mediate bidirectional transfer of signals across the plasma membrane. The cytoplasmic domains of integrin family members interact with components of the signal transduction apparatus within cells. Integrin receptors contain noncovalently associated α and β subunits that consist of a large extracellular region (the ligand-binding domain), a short transmembrane region, and a cytoplasmic domain of varying length. In mammals, at least 17 α subunits and 8 β subunits have been identified and these proteins can heterodimerize to form at least 22 different receptors. The integrin β 2 subunit associates with integrin α L to form a receptor for ICAM family members. Integrin β 2/ α L is involved in a variety of immune phenomena including leukocyte-endothelial cell interaction, cytotoxic T-cell mediated killing, and antibody dependent killing by granulocytes and monocytes.

Note:

Protein G purified tissue culture supernatant.