

Product datasheet for TA389215

STAT1 Mouse Antibody [Clone ID: M249]

Product data:

Product Type: Primary Antibodies

Clone Name: M249

Applications: ICC, IP, WB Recommended Dilution: **WB**: 1:1000

ICC: 1:100

Reactivity: Human, Rat, Mouse

Host: Mouse Isotype: IgG1

Immunogen: Clone (M249) was generated from a recombinant protein from human Stat1. This sequence

has high homology to the conserved region in rat and mouse stat1 a and b.

Specificity: The antibody detects 84 and 91 kDa* stat1 variants on SDS-PAGE immunoblots of human

Jurkat and A431 cells.

Formulation: PBS + 1 mg/ml BSA, 0.05% NaN3 and 50% glycerol

Concentration: lot specific

Purification: Protein A Purified

Conjugation: Unconjugated

Storage: Storage at -20°C is recommended, as aliquots may be taken without freeze/thawing due to

presence of 50% glycerol. Stable for at least 1 year at -20°C.

Stability: After date of receipt, stable for at least 1 year at -20°C.

Predicted Protein Size: 84/91

Database Link: P42224



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



STAT1 Mouse Antibody [Clone ID: M249] - TA389215

Background:

The stat proteins function both as cytoplasmic signal transducers and as activators of transcription. Stat1 is expressed as two variants of 84 and 91 kDa. Stat1 proteins contain SH2 and SH3 domains, and are components of the interferon-stimulated gene factor 3 (ISGF3) complex. This complex is the primary transcription activator induced by the binding of interferon to its receptors. In response to activation by various cytokines and growth factors, stat1 subunits become phosphorylated at tyrosine 701. This leads to translocation of stat1 to the nucleus, resulting in formation of an active ISGF3 complex. Active ISGF3 modulates the transcription of the interferon-stimulated genes. Thus, phosphorylation of Tyr-701 is critical for gene expression mediated by various cytokines and growth factors.

Note:

Protein G purified tissue culture supernatant.