

Product datasheet for TA389300

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Biotinylated Anti-CD138 antibody(DM56), Rabbit mAb

Product data:

Product Type: Primary Antibodies

Clone Name: DM56

Applications: ELISA, FC

Recommended Dilution: ELISA 1:5000-10000

Flow Cyt 1:100

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Monoclonal

Formulation: 1XPBS

Concentration: Lot specific

Purification: Purified from cell culture supernatant by affinity chromatography

Conjugation: Unconjugated

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and

thawing).Lyophilized antibodies are shipped at ambient temperature.

Stability: 12 months from date of despatch

Background: Syndecan-1 (SYND1 or SDC1) is also known as CD antigen CD138, is a transmembrane (type I)

heparan sulfate proteoglycan and is a member of the syndecan proteoglycan family. The syndecans mediate cell binding, cell signaling, and cytoskeletal organization and syndecan receptors are required for internalization of the HIV-1 tat protein. The syndecan-1 / SDC1 protein functions as an integral membrane protein and participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. It is a useful marker for plasma cells, but only if the cells tested are already known to be derived

from blood.

Synonyms: SDC1, Syndecan-1, CD138, SYND1, SDC

