

## **Product datasheet for TA389467**

## 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com

OriGene Technologies, Inc.

techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Biotinylated Anti-CTLA4 antibody(DM51), Rabbit mAb

## **Product data:**

**Product Type:** Primary Antibodies

Clone Name: DM51

**Applications:** ELISA, FC

Recommended Dilution: ELISA 1:5000-10000

Flow Cyt 1:100

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Monoclonal

Formulation: 1XPBS

**Concentration:** Lot specific

**Purification:** Purified from cell culture supernatant by affinity chromatography

**Conjugation:** Unconjugated

Storage: Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended

for use within a month, aliquot and store at -80°C (Avoid repeated freezing and

thawing).Lyophilized antibodies are shipped at ambient temperature.

Stability: 12 months from date of despatch

Background: Leukocyte surface antigen CD47 is also known as Antigenic surface determinant protein OA3,

Integrin-associated protein (IAP) and Protein MER6. CD47 contains 1 Ig-like V-type (immunoglobulin-like) domain. CD47 is very broadly distributed on normal adult tissues. CD47 has a role in both cell adhesion by acting as an adhesion receptor for THBS1 on platelets, and in the modulation of integrins and plays an important role in memory

formation and synaptic plasticity in the hippocampus by similarity. CD47 is the receptor for SIRPA, binding to which prevents maturation of immature dendritic cells and inhibits cytokine production by mature dendritic cells. CD47 Interaction with SIRPG mediates cell-cell adhesion, enhances superantigen-dependent T-cell-mediated proliferation and costimulates T-cell

ennances superantigen-dependent 1-cen-mediated promeration and costimulates 1-cen

activation.

Synonyms: CTLA4, CD152

