

Product datasheet for TA389547

Anti-CCR2 antibody(DMC447), IgG1 Chimeric mAb

Product data:

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

| Product Type: | Primary Antibodies |
|-----------------------|---|
| Applications: | FC |
| Recommended Dilution: | Flow Cyt 1:100 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Monoclonal |
| Formulation: | 1XPBS |
| Concentration: | Lot specific |
| Purification: | Purified from cell culture supernatant by affinity chromatography |
| Conjugation: | Unconjugated |
| Storage: | Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized antibodies are shipped at ambient temperature. |
| Stability: | 12 months from date of despatch |
| Background: | The protein encoded by this gene is a receptor for monocyte chemoattractant protein-1, a chemokine which specifically mediates monocyte chemotaxis. Monocyte chemoattractant protein-1 is involved in monocyte infiltration in inflammatory diseases such as rheumatoid arthritis as well as in the inflammatory response against tumors. The encoded protein mediates agonist-dependent calcium mobilization and inhibition of adenylyl cyclase. This protein can also be a coreceptor with CD4 for HIV-1 infection. This gene is located in the chemokine receptor gene cluster region of chromosome 3. |
| Synonyms: | CC-CKR-2, MCP-1-R, CD192 |



This product is to be used for laboratory only. Not for diagnostic or therapeutic use. ©2023 OriGene Technologies, Inc., 9620 Medical Center Drive, Ste 200, Rockville, MD 20850, US