

## Product datasheet for **TA389564**

### Anti-CD164 antibody(DMC476), IgG1 Chimeric mAb

#### Product data:

Product Type:	Primary Antibodies
Applications:	FC
Recommended Dilution:	Flow Cyt 1:100
Reactivity:	Human
Host:	Rabbit
Clonality:	Monoclonal
Formulation:	1XPBS
Concentration:	Lot specific
Purification:	Purified from cell culture supernatant by affinity chromatography
Conjugation:	Unconjugated
Storage:	Store at -20°C to -80°C for 12 months in lyophilized form. After reconstitution, if not intended for use within a month, aliquot and store at -80°C (Avoid repeated freezing and thawing).Lyophilized antibodies are shipped at ambient temperature.
Stability:	12 months from date of despatch
Background:	This gene encodes one of at least three opioid receptors in humans; the mu opioid receptor (MOR). The MOR is the principal target of endogenous opioid peptides and opioid analgesic agents such as beta-endorphin and enkephalins. The MOR also has an important role in dependence to other drugs of abuse, such as nicotine, cocaine, and alcohol via its modulation of the dopamine system. The NM_001008503.2:c.118A>G allele has been associated with opioid and alcohol addiction and variations in pain sensitivity but evidence for it having a causal role is conflicting. Multiple transcript variants encoding different isoforms have been found for this gene. Though the canonical MOR belongs to the superfamily of 7-transmembrane-spanning G-protein-coupled receptors some isoforms of this gene have only 6 transmembrane domains. [provided by RefSeq, Oct 2013]
Synonyms:	LMOR; M-OR-1; MOP; MOR; MOR1; OPRM



[View online »](#)