

## **Product datasheet for TA389681**

## **XBP1S Rabbit Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

**Applications:** ICC/IF, WB

Recommended Dilution: WB,1:100 - 1:500

IF/ICC,1:50 - 1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Formulation:** PBS with 0.05% proclin300,50% glycerol,pH7.3.

**Concentration:** lot specific

**Conjugation:** Unconjugated

Storage: Store at -20°C as received.

**Stability:** Stable for 12 months from date of receipt.

Database Link: P17861-2

Background: This gene encodes a transcription factor that regulates MHC class II genes by binding to a

promoter element referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRNA of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XBP1(S), which is the functionally active transcription factor. The isoform encoded by the unspliced mRNA, XBP1(U), is constitutively expressed, and thought to function as a negative feedback regulator of XBP1(S), which shuts off transcription of target genes during the recovery phase of ER stress. A pseudogene of

XBP1 has been identified and localized to chromosome 5.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com