

Product datasheet for TA392481M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Poliovirus Receptor (PVR) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 1:5000~1:10000

Reactivity: Human, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to Human CD155.

Specificity: CD155 polyclonal antibody detects endogenous levels of CD155 protein. **Formulation:** Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 70 kDa

Gene Name: poliovirus receptor

Database Link: Entrez Gene 5817 Human

P15151

Background: Poliovirus receptor (PVR, CD155) is an immunoglobulin-like, transmembrane glycoprotein

originally described as a mediator of poliovirus attachment to cells and later identified as important in adherens junction formation. Also known as nectin-like 5 (Necl-5), PVR binds nectin-3 and interacts with integrin $\alpha\nu\beta$ 3 and PDGFR to regulate integrin clustering and focal contact formation at the leading edge of migrating cells. Research studies demonstrate that

PVR and nectin-3 regulate contact inhibition during cell motility and proliferation in

transformed 3T3 cells. Additional research indicates that PVR (CD155, Necl-5) expression may play a role in invasiveness of lung adenocarcinoma. In the immune system, CD155 plays a

role in natural killer (NK) cell-mediated cytotoxicity.

Synonyms: CD155; Nectin-like protein 5 (NECL-5); Poliovirus receptor; PVR; PVS






Note:

For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of CD155 polyclonal antibody at 1:10000 dilution Lane1:CT-26 whole cell lysate(30ug) Lane2:HepG2 whole cell lysate(10ug) Lane3:A549 whole cell lysate(10ug)