

Product datasheet for TA392507

IRS1 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: IF, WB

Recommended Dilution: WB: 1:10000~1:20000 IF 1:50 - 1:100

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to Human IRS-1.

Specificity: IRS-1 (S1101) polyclonal antibody detects endogenous levels of IRS-1 protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2.

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 190 kDa

Gene Name: insulin receptor substrate 1

Database Link: Entrez Gene 3667 Human

P35568

Background: Insulin receptor substrate 1 (IRS-1) is one of the major substrates of the insulin receptor

kinase. IRS-1 contains multiple tyrosine phosphorylation motifs that serve as docking sites for SH2-domain containing proteins that mediate the metabolic and growth-promoting functions

of insulin. IRS-1 also contains over 30 potential serine/threonine phosphorylation sites. Ser307 of IRS-1 is phosphorylated by JNK and IKK while Ser789 is phosphorylated by SIK-2, a member of the AMPK family. The PKC and mTOR pathways mediate phosphorylation of IRS-1 at Ser612 and Ser636/639, respectively. Phosphorylation of IRS-1 at Ser1101 is mediated by PKCθ and results in an inhibition of insulin signaling in the cell, suggesting a potential

mechanism for insulin resistance in some models of obesity.

Synonyms: Insulin receptor substrate 1; IRS-1; IRS1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Note:

For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of IRS-1 (S1101) polyclonal antibody at 1:10000 dilution Lane1:HepG2 whole cell lysate(40ug) Lane2:MCF-7 whole cell lysate(20ug)



Immunofluorescence analysis of MCF-7 cells using IRS-1 antibody at dilution of 1:100.