

# **Product datasheet for TA392622S**

### **OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Cytochrome C (CYCS) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Recommended Dilution:** WB: 1:2000~1:5000 IHC:1:50~1:100

**Reactivity:** Human, Rat, Mouse

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant protein of human Cytochrome c.

**Specificity:** Cytochrome c polyclonal antibody detects endogenous levels of Cytochrome c protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.4.

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 17 kDa

Gene Name: cytochrome c, somatic

Database Link: Entrez Gene 54205 Human

P99999

**Background:** Cytochrome c is a well conserved electron-transport protein and is part of the respiratory

chain localized to mitochondrial intermembrane space. Upon apoptotic stimulation,

cytochrome c released from mitochondria associates with procaspase-9 (47 kDa)/Apaf 1. This complex processes caspase-9 from inactive proenzyme to its active form. This event further

triggers caspase-3 activation and eventually leads to apoptosis.


**Synonyms:** CYC; CYCS; Cytochrome c

**Note:** For research use only, not for use in diagnostic procedure.






# **Product images:**



Immunohistochemistry of paraffin-embedded human colon carcinoma using Cytochrome c antibody at dilution of 1:50.



Immunohistochemistry of paraffin-embedded human colon carcinoma using Cytochrome c antibody at dilution of 1:50.