

## **Product datasheet for TA392678S**

## COX4 (COX4I1) Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WB

**Reactivity:** WB: 1:500~1:1000 Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Recombinant full length Human Cox-4.

**Specificity:** Cox-4 polyclonal antibody detects endogenous levels of Cox-4 protein. **Formulation:** Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 20 kDa

**Gene Name:** cytochrome c oxidase subunit 4l1

**Database Link:** Entrez Gene 1327 Human

P13073

**Background:** Cytochrome c oxidase (COX) functions as the terminal oxidase of the respiratory chain that

uses cytochrome c as an electron donor to drive a proton gradient across the inner mitochondrial membrane. The mammalian COX apoenzyme is a heteromer consisting of three mitochondrial encoded catalytic subunits and several nuclear gene encoded structural subunits. COX contains two iron-coordination sites and two copper-coordination sites.

Cytochrome c oxidase IV (COX4) is a nuclear-encoded subunit of COX that may play a role in regulating COX activity. COX4 is expressed ubiquitously in adult human tissue with the strongest levels of expression in the pancreas and moderate expression levels in heart,

skeletal muscle and placenta.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: COX4; COX4I1; COX IV-1; Cytochrome c oxidase polypeptide IV; Cytochrome c oxidase subunit

4 isoform 1, mitochondrial; Cytochrome c oxidase subunit IV isoform 1

**Note:** For research use only, not for use in diagnostic procedure.

## **Product images:**



Western blot (WB) analysis of Cox-4 pAb at 1:500 dilution Lane1:3T3-L1 whole cell lysate(40ug) Lane2:C6 whole cell lysate(40ug) Lane3:HCT116 whole cell lysate(10ug) Lane4:U-87MG whole cell lysate(10ug)