

Product datasheet for TA392708S

Product data:

Product Type: Primary Antibodies

PDPK1 Rabbit Polyclonal Antibody

Applications:

Recommended Dilution: WB: 1:500~1:1000 IHC: 1:50~1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 211-260 of Human PDPK1.

Specificity: PDPK1 (K235) polyclonal antibody detects endogenous levels of PDPK1 protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles. Storage:

Stability: 1 year **Predicted Protein Size:** ~ 63 kDa

Gene Name: 3-phosphoinositide dependent protein kinase 1

Database Link: Entrez Gene 5170 Human

O15530

Background: PDK1 (3 Phosphoinositide Dependent Protein Kinase 1) phosphorylates AGC kinases. PDK1

activates conventional PKC and PKC zeta through phosphorylation of critical threonine

residues in the activation loop. PDK1 also phosphorylates Protein Kinase B (PKB) at threonine

308 in the presence of hosphatidylinositol-3,4,5-trisphosphate. Active Akt inactivates Glycogen Synthase Kinase 3 (GSK3), eventually leading to the dephosphorylation and activation of glycogen synthase, and the stimulation of glycogen synthesis. Because of the role that PDK plays in insulin-induced glycogen synthesis and PKC activation, it is a potentially

important target for metabolic drug research.

Synonyms: 3-phosphoinositide-dependent protein kinase 1; hPDK1; PDK1; PDPK1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Note:

For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of PDPK1 (K235) Antibody at 1:1000 dilution Lane1:MCF-7 whole cell lysate(40ug) Lane2:A549 whole cell lysate(40ug) Lane3:PC12 whole cell lysate(40ug) Lane4:3T3-L1 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of PDPK1 (K235) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.