

Product datasheet for TA392732

FRA1 (FOSL1) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500~1:1000 IHC: 1:50~1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 100-150 of Human Fra-1.

Specificity: Fra-1 (D134) polyclonal antibody detects endogenous levels of Fra-1 protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 30, 42 kDa

Gene Name: FOS like 1, AP-1 transcription factor subunit

Database Link: Entrez Gene 8061 Human

P15407

Background: The v-Fos oncogene was initially detected in two independent murine osteosarcoma virus

isolates and an avian nephroblastoma virus. Members of the c-Fos gene family, including c-Fos, Fos B, Fra-1 and Fra-2, encode nuclear phosphoproteins that are rapidly and transiently induced by a variety of agents and function as transcriptional regulators for several genes. In contrast to c-Jun proteins, which form homo- and heterodimers that bind to specific DNA response elements, c-Fos proteins are only active as heterodimers with members of the Jun gene family. In addition, selected ATF/CREB family members can form leucine zipper dimers with Fos and Jun. Different dimers exhibit differential specificity and affinity for AP-1 and CRE

sites.

Synonyms: Fos-related antigen 1; FOSL1; FRA-1; FRA1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Note:

For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of Fra-1 (D134) pAb at 1:500 dilution Lane1:PC3 whole cell lysate(40ug) Lane2:HepG2 whole cell lysate(40ug) Lane3:The Brain tissue lysate of Mouse(40ug) Lane4:PC12 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of Fra-1 (D134) pAb in paraffin-embedded human breast carcinoma tissue at 1:50.