

Product datasheet for TA392830S

ABCA8 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB: 1:500~1:1000 Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 1171-1220 of Human ABCA8. Specificity: ABCA8 (F1195) polyclonal antibody detects endogenous levels of ABCA8 protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles. Storage:

Stability: 1 year

Predicted Protein Size: ~ 179 kDa

Gene Name: ATP binding cassette subfamily A member 8

Database Link: Entrez Gene 10351 Human

O94911

Background: ATP-binding cassette (ABC) transporters are an evolutionarily conserved family of proteins

> that catalyze the transport of molecules across extracellular and intracellular membranes by harnessing the energy of ATP hydrolysis. ABCA8 (ATP-binding cassette, sub-family A (ABC1), member 8) is a 1,581 amino acid multi-pass membrane protein that contains two ABC transporter domains and belongs to the ABC transporter family. Expressed in a variety of tissues with particularly high expression in liver, heart and skeletal muscle, ABCA8 functions as an ATP-dependent lipophilic drug transporter that exists as multiple alternatively spliced

isoforms.

Synonyms: ATP-binding cassette sub-family A member 8; KIAA0822



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Note:

For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of ABCA8 (F1195) pAb at 1:500 dilution Lane1:H9C2 whole cell lysate(40ug) Lane2:AML-12 whole cell lysate(40ug) Lane3:SK-OVCAR3 whole cell lysate(40ug) Lane4:HepG2 whole cell lysate(40ug) Lane5:L02 whole cell lysate(10ug)