

## Product datasheet for TA392831

## **DQX1 Rabbit Polyclonal Antibody**

**Product data:** 

**Product Type: Primary Antibodies** 

**Applications:** 

Recommended Dilution: WB: 1:500~1:1000 Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 560-610 of Human DQX1.

Specificity: DQX1 (L595) polyclonal antibody detects endogenous levels of DQX1 protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles. Storage:

Stability: 1 year **Predicted Protein Size:** ~ 66 kDa

Gene Name: DEAQ-box RNA dependent ATPase 1

Entrez Gene 165545 Human Database Link:

Q8TE96

Background: DQX1 (DEAQ box RNA-dependent ATPase 1), also known as FLJ23757, is a 71 amino acid

> protein that contains one helicase ATP-binding domain and one helicase C-terminal domain. Localized to the nucleus, DQX1 catalyzes the conversion of ATP to ADP and a phosphate. Expressed as three isoforms produced by alternative splicing events, DQX1 is encoded by a gene that maps to human chomosome 2. As the second largest human chromosome, chromosome 2 makes up approximately 8% of the human genome and contains 237 million

bases encoding over 1,400 genes. A number of genetic diseases are linked to genes on chromosome 2, including Harlequin icthyosis, sitosterolemia and Alström syndrome.

Synonyms: ATP-dependent RNA helicase DQX1; DEAQ box polypeptide 1



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com



Note:

For research use only, not for use in diagnostic procedure.

## **Product images:**



Western blot (WB) analysis of DQX1 (L595) pAb at 1:500 dilution Lane1:HepG2 whole cell lysate(40ug) Lane2:HCT116 whole cell lysate(40ug) Lane3:SGC7901 whole cell lysate(40ug) Lane4:CT26 whole cell lysate(40ug) Lane5:H9C2 whole cell lysate(40ug)