

Product datasheet for TA392835

MED18 Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB: 1:500~1:1000 Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Immunogen: Synthetic peptide, corresponding to amino acids 72-120 of Human MED18.

Specificity: MED18 (V102) polyclonal antibody detects endogenous levels of MED18 protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles. Storage:

Stability: 1 year **Predicted Protein Size:** ~ 24 kDa

Gene Name: mediator complex subunit 18

Database Link: Entrez Gene 54797 Human

Q9BUE0

Background: In mammalian cells, transcription is regulated in part by high molecular weight co-activating

> complexes that mediate signals between transcriptional activators and RNA polymerase II (Pol II). The mediator complex is one such multi-protein structure that functions as a bridge between regulatory proteins and Pol II, thereby regulating Pol II-dependent transcription. Med18 (mediator complex subunit 18), also known asp28b, is a 208 amino acid protein that localizes to nucleus and exists as a component of the mediator complex. Working in tandem with several other proteins, including Med8 and Med25, Med18 serves as a scaffold for the assembly of a functional preinitiation complex with Pol II and general transcription factors,

thereby activating the transcription of Pol II-dependent genes.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms: MED18; Mediator complex subunit 18; Mediator of RNA polymerase II transcription subunit

18; p28b

Note: For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of MED18 (V102) pAb at 1:1000 dilution Lane1:HepG2 whole cell lysate(20ug) Lane2:SGC7901 whole cell lysate(20ug) Lane3:The Brain tissue lysate of Rat(40ug) Lane4:The Brain tissue lysate of Mouse(40ug)