

## **Product datasheet for TA392848**

## CDC40 Rabbit Polyclonal Antibody

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Recommended Dilution:** WB: 1:500~1:1000 IHC: 1:50~1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

**Immunogen:** Synthetic peptide, corresponding to amino acids 181-230 of Human Cdc40.

**Specificity:** Cdc40 (S209) polyclonal antibody detects endogenous levels of Cdc40 protein.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

**Storage:** Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 65 kDa

**Gene Name:** cell division cycle 40

**Database Link:** Entrez Gene 51362 Human

<u>060508</u>

**Background:** Cdc40, also known as pre-mRNA-processing factor 17 (PRPF17) or EH-binding protein 3

(EHB3), is a 579 amino acid nuclear protein. Cdc40 is essential for the catalytic step II of the pre-mRNA splicing process, in which Cdc40 associates with the spliceosome C complex. Cdc40 contains seven WD repeats, which are important in protein-protein interactions. Cdc40

has sequence similarity to the yeast protein Prp17, which is involved in pre-mRNA splicing and cell cycle progression. The sequence similarity between the mammalian Cdc40 and the yeast Prp17 may indicate an additional role in cell cycle progression for mammalian Cdc40.

Synonyms: CDC40; Cell division cycle 40 homolog; EH-binding protein 3; EHB3; Ehb3; hPRP17; Pre-mRNA-

processing factor 17; PRP17; PRP17 homolog; PRPF17



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Note:

For research use only, not for use in diagnostic procedure.

## **Product images:**



Western blot (WB) analysis of Cdc40 (S209) pAb at 1:500 dilution Lane1:HEK293T whole cell lysate(10ug) Lane2:MCF-7 whole cell lysate(10ug) Lane3:CT26 whole cell lysate(10ug) Lane4:The Brain tissue lysate of Rat(40ug)



Immunohistochemistry (IHC) analyzes of Cdc40 (S209) pAb in paraffin-embedded human colorectal carcinoma tissue at 1:50.