

Product datasheet for **TA392867S**

ACAT1 Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:500~1:1000 IHC: 1:50~1:200
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic peptide, corresponding to amino acids 235-286 of Human ACAT1.
Specificity:	ACAT1 (K266) polyclonal antibody detects endogenous levels of ACAT1 protein.
Formulation:	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Concentration:	1mg/ml
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.
Stability:	1 year
Predicted Protein Size:	~ 45 kDa
Gene Name:	acetyl-CoA acetyltransferase 1
Database Link:	Entrez Gene 38 Human P24752
Background:	ACAT-1 (acetyl-Coenzyme A acetyltransferase 1) is a mitochondrial enzyme involved in the formation and degradation of ketone bodies and is necessary for the proper metabolic processing of isoleucine. Rare defects in the gene encoding ACAT-1 lead to β -ketothiolase deficiency, which is characterized by ketoacidotic attacks. ACAT-2 (acetyl-Coenzyme A acetyltransferase 2) is considered a cytosolic protein and is crucial for cholesterol synthesis. Specifically, both Acetoacetyl-CoA specific thiolases, ACAT-1 and ACAT-2 catalyze the formation of acetoacetyl-CoA from two acetyl-CoA molecules. These enzymes are also capable of the reverse reaction, the cleavage of acetoacetyl-CoA into two acetyl-CoA molecules.




[View online »](#)


Synonyms: ACAT; ACAT-1; ACAT1; Acetoacetyl-CoA thiolase; Acetyl-CoA acetyltransferase, mitochondrial; MAT

Note: For research use only, not for use in diagnostic procedure.

Product images:



Immunohistochemistry (IHC) analyzes of ACAT1 (K266) pA bin paraffin-embedded human liver carcinoma tissue at 1:50, showing cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.



Immunohistochemistry (IHC) analyzes of ACAT1 (K266) pA bin paraffin-embedded human kidney carcinoma tissue at 1:50, showing cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.