

Product datasheet for TA392885

CRKL Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB: 1:500~1:1000 IHC: 1:50~1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Synthetic phosphopeptide derived from human CrkL around the phosphorylation site of Immunogen:

Tyrosine 207.

Specificity: p-Crk-L (Y207) polyclonal antibody detects endogenous levels of Crk-L protein only when

phosphorylated at Tyr207.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year **Predicted Protein Size:** ~ 39 kDa

Gene Name: v-crk avian sarcoma virus CT10 oncogene homolog-like

Database Link: Entrez Gene 1399 Human

P46109

Background: Crk, is a transformation-specific protein that induces elevation of cellular phosphotyrosine

> levels, but lacks tyrosine kinase activity itself. A second protein, Nck, consists solely of three SH3 domains and one SH2 domain, while GRB2 contains an SH2 domain flanked on both sides by SH3 domains. A member of this protein class, Crk-L, is encoded by a gene located on chromosome 22, band 11, centromeric of the chronic myelogenous leukemia breakpoint region. Crk-L encodes a 303 amino acid protein with one SH2 and two SH3 domains, and a

predicted molecular weight of 36 kDa.

Synonyms: Crk-like protein; Crk L; Crkl; Crk like protein; Crkol; Oncogene CrkL



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn


Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Note:

For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of p-CrkL (Y207) pAb at 1:500 dilution Lane1:MCF-7 whole cell lysate(40ug) Lane2:HEK293T whole cell lysate(40ug) Lane3:PMVEC whole cell lysate(40ug) Lane4:CT26 whole cell lysate(40ug) Lane5:SGC7901 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of p-Crk-L (Y207) pAb in paraffin-embedded human colon cancer tissue.