

Product datasheet for TA392896M

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Serum Response Factor (SRF) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WE

Recommended Dilution: WB: 1:500~1:1000 IHC: 1:50~1:200

Reactivity: Human
Host: Rabbit
Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic phosphopeptide derived from human SRF around the phosphorylation site of

Serine 103.

Specificity: p-SRF (S103) polyclonal antibody detects endogenous levels of SRF protein when

phosphorylated at Ser103.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year Predicted Protein Size: ~ 52 kDa

Gene Name: serum response factor

Database Link: Entrez Gene 6722 Human

P11831






Background:

Serum response factor (SRF) is a transcription factor that binds the serum response element (SRE), a sequence that mediates the transient response of many cellular genes to growth stimulation. SRF-binding sites are also constitutive promotor elements in many muscle-specific promotors. At the c-Fos SRE, formation of a ternary complex containing SRF and its accessory protein p62TCF appears to be important for signal transduction. Two related Ets domain proteins, Elk-1 and SRF accessory protein-1 (SAP-1) have DNA binding properties identical to that of p62TCF. Elk-1 and SAP-1 contain two homologous regions of which the two amino-terminal regions, the Ets domain (box A) and the B box, mediate ternary complex formation with SRF.


Synonyms: Serum response factor

Note: For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of SRF (phospho-S103) polyclonal antibody at 1:500 dilution Lane1:H1792 whole cell lysate(40ug) Lane2:SK-OVCAR3 whole cell lysate(40ug) Lane3:HEK293T whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of p-SRF (S103) pAb in paraffin-embedded human lung cancer tissue.