

Product datasheet for TA392907M

POLR2A Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications:

Recommended Dilution: WB: 1:500~1:1000 IHC: 1:50~1:200

Reactivity: Human, Mouse, Rat

Host: Rabbit Isotype: lgG

Clonality: Polyclonal

Synthetic phosphopeptide derived from human Rpb1 around the phosphorylation site of Immunogen:

Serine 1619.

Specificity: p-Rpb1 (S1619) polyclonal antibody detects endogenous levels of Rpb1 protein only when

phosphorylated at Ser1619.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year **Predicted Protein Size:** ~ 280 kDa

Gene Name: polymerase (RNA) II subunit A

Database Link: Entrez Gene 5430 Human

P24928

Background: RPB1 is a subunit of RNA polymerase II which catalyses transcription of RNA from DNA. RPB1

> forms the polymerase active center together with the second largest subunit. Pol II is the central component of the basal RNA polymerase II transcription machinery. It is composed of mobile elements that move relative to each other. RPB1 is part of the core element with the central large cleft, the clamp element that moves to open and close the cleft and the jaws that

are thought to grab the incoming DNA template.



OriGene Technologies, Inc. 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

DNA-directed RNA polymerase III largest subunit; DNA-directed RNA polymerase II subunit A; DNA-directed RNA polymerase II subunit RPB1; POLR2; POLR2A; RNA-directed RNA polymerase II subunit RPB1; RNA polymerase II subunit B1


Note:

For research use only, not for use in diagnostic procedure.

Product images:



Western blot (WB) analysis of p-Rpb1 (S1619) pAb at 1:2000 dilution Lane1:A549 whole cell lysate(40ug) Lane2:Hela whole cell lysate(40ug) Lane3:A2780 whole cell lysate(40ug) Lane4:PC12 whole cell lysate(40ug) Lane5:3T3-L1 whole cell lysate(40ug)



Immunohistochemistry (IHC) analyzes of p-Rpb1 (S1619) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.