

### **Product datasheet for TA392937S**

#### OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

## Tyrosine Hydroxylase (TH) Rabbit Polyclonal Antibody

#### **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

Recommended Dilution: WB: 1:500~1:1000

Reactivity: Human, Mouse, Rat

**Host:** Rabbit

**Isotype:** IgG

**Clonality:** Polyclonal

**Immunogen:** Synthetic peptide derived from human TH around the phosphorylation site of S8.

Specificity: TH (phospho-S8) polyclonal antibody detects endogenous levels of TH protein only when

phosphorylated at Ser8.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

**Conjugation:** Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 58 kDa

**Gene Name:** tyrosine hydroxylase

**Database Link:** Entrez Gene 7054 Human

P07101

Background: The enzyme tyrosine hydroxylase (TH), also designated tyrosine 3-monooxygenase (TY3H),

catalyzes the conversion of tyrosine to L-dopa, which is the rate limiting step in the

biosynthesis of catecholamines such as dopamine, adrenalin and noradrenalin. TH is thought to play a role in the pathogenesis of Parkinson's disease, which is associated with reduced dopamine levels. Two transcription factor binding sites in the proximal region of the TH gene, the TPA-responsive element (TRE) and the c-AMP responsive element (CRE), have been implicated in the complex regulation of the TH gene. TH is also known to be upregulated by

the glia maturation factor (GMF), a Cdc 10/SWI6 motif-containing protein called V-1, and a

variety of additional compounds






**Synonyms:** TH; TYH; Tyrosine 3-hydroxylase; Tyrosine 3-monooxygenase

**Note:** For research use only, not for use in diagnostic procedure.

# **Product images:**



Western blot (WB) analysis of p-TH (S8) pAb at 1:500 dilution Lane1:MCF-7 whole cell lysate(40ug) Lane2:MCF-7 treated with Forskolin(4µg/ml,30 minutes) whole cell lysate(40ug) Lane3:MCF-7 treated with Forskolin(4µg/ml,60 minutes) whole cell lysate(40ug)