

Product datasheet for **TA392944S**

Vimentin (VIM) Rabbit Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	WB
Recommended Dilution:	WB: 1:500~1:1000
Reactivity:	Human, Mouse, Rat
Host:	Rabbit
Isotype:	IgG
Clonality:	Polyclonal
Immunogen:	Synthetic phosphopeptide derived from human Vimentin around the phosphorylation site of Serine 56.
Specificity:	p-Vimentin (S56) polyclonal antibody detects endogenous levels of Vimentin protein when phosphorylated at Ser56.
Formulation:	Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2
Concentration:	1mg/ml
Conjugation:	Unconjugated
Storage:	Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.
Stability:	1 year
Predicted Protein Size:	~ 40, 57 kDa
Gene Name:	vimentin
Database Link:	Entrez Gene 7431 Human P08670



[View online »](#)

Background:


Phosphorylation of Vimentin induces disassembly of Vimentin intermediate filaments in vivo and in vitro. Binding of 14-3-3 depends on Vimentin phosphorylation and requires the phosphopeptide binding domain of 14-3-3, which is an amino terminal head domain consisting of amino acids 1-96. Phosphorylated Vimentin sequesters 14-3-3 and limits its availability to other target proteins, which can affect intracellular signaling processes that require 14-3-3. The amino-terminal domain of Vimentin is the target site for several protein kinases, including Rho kinase and PKC. Ser 38 and Ser 71 of Vimentin are the major sites of phosphorylation by Rho kinase. The disruption of subcellular compartmentalization of interphase cells leads to PKC-mediated phosphorylation of Vimentin. Thus, targeting of activated PKC, coupled with the reorganization of intracellular membranes, which contain phospholipids essential for activation, leads to the mitosis-specific phosphorylation of Vimentin.

Synonyms:

VIM; Vimentin

Note:

For research use only, not for use in diagnostic procedure.

Product images:

Western blot (WB) analysis of p-Vimentin (S56) pAb at 1:500 dilution Lane 1: MCF-7 whole cell lysate (40µg) Lane 2: A375 whole cell lysate (40µg) Lane 3: C6 whole cell lysate (40µg) Lane 4: The testis tissue lysate of Mouse (40µg) Lane 5: HeLa whole cell lysate (40µg)