

Product datasheet for **TA392946S**

14-3-3 theta (YWHAQ) Rabbit Polyclonal Antibody

Product data:

| | |
|-------------------------|---|
| Product Type: | Primary Antibodies |
| Applications: | WB |
| Recommended Dilution: | WB: 1:500~1:1000 |
| Reactivity: | Human, Mouse, Rat |
| Host: | Rabbit |
| Isotype: | IgG |
| Clonality: | Polyclonal |
| Immunogen: | Synthetic phosphopeptide derived from human 14-3-3 θ/τ around the phosphorylation site of Serine 232 |
| Specificity: | 14-3-3 θ/τ (phospho-S232) polyclonal antibody detects endogenous levels of 14-3-3 θ/τ protein only when phosphorylated at Ser232. |
| Formulation: | Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2 |
| Concentration: | 1mg/ml |
| Conjugation: | Unconjugated |
| Storage: | Store at 4°C short term. Aliquot and store at -22°C long term. Avoid freeze-thaw cycles. |
| Stability: | 1 year |
| Predicted Protein Size: | ~ 28 kDa |
| Gene Name: | tyrosine 3-monooxygenase/tryptophan 5-monooxygenase activation protein theta |
| Database Link: | Entrez Gene 10971 Human P27348 |



[View online »](#)

Background:


The 14-3-3 family of proteins plays a key regulatory role in signal transduction, checkpoint control, apoptotic and nutrient-sensing pathways. 14-3-3 proteins are highly conserved and ubiquitously expressed. There are at least seven isoforms, β , γ , ϵ , σ , ζ , τ , and η that have been identified in mammals. The initially described α and δ isoforms are confirmed to be phosphorylated forms of β and ζ , respectively. Through their amino-terminal α helical region, 14-3-3 proteins form homo- or heterodimers that interact with a wide variety of proteins: transcription factors, metabolic enzymes, cytoskeletal proteins, kinases, phosphatases, and other signaling molecules. The interaction of 14-3-3 proteins with their targets is primarily through a phospho-Ser/Thr motif. However, binding to divergent phospho-Ser/Thr motifs, as well as phosphorylation independent interactions has been observed. 14-3-3 binding masks specific sequences of the target protein, and therefore, modulates target protein localization, phosphorylation state, stability, and molecular interactions. 14-3-3 proteins may also induce target protein conformational changes that modify target protein function. Distinct temporal and spatial expression patterns of 14-3-3 isoforms have been observed in development and in acute response to extracellular signals and drugs, suggesting that 14-3-3 isoforms may perform different functions despite their sequence similarities. Several studies suggest that 14-3-3 isoforms are differentially regulated in cancer and neurological syndromes.

Synonyms:

14-3-3 protein T-cell; 14-3-3 protein tau; 14-3-3 protein theta; Protein HS1; YWHAQ

Note:

For research use only, not for use in diagnostic procedure.

Product images:

Western blot (WB) analysis of 14-3-3 θ/τ (phospho-S232) polyclonal antibody at 1:500 dilution Lane1:A549 whole cell lysate(40 μ g) Lane2:Hela whole cell lysate(40 μ g) Lane3:The Brain tissue lysate of Rat(40 μ g) Lane4:HEK293T whole cell lysate(40 μ g) Lane5:CT-26 whole cell lysate(40 μ g)