

Product datasheet for TA392949

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

Cytokeratin 18 (KRT18) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Recommended Dilution: WB: 1:500~1:1000

Reactivity: Human

Host: Rabbit

Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic phosphopeptide derived from human Cytokeratin 18 around the phosphorylation

site of Serine 52

Specificity: p-Cytokeratin 18 (S52) polyclonal antibody detects endogenous levels of Cytokeratin 18

protein only when phosphorylated at Ser52.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 46 kDa

Gene Name: keratin 18

Database Link: Entrez Gene 3875 Human

P05783

Background: Keratins (cytokeratins) are intermediate filament proteins that are mainly expressed in

epithelial cells. Keratin heterodimers composed of an acidic keratin (or type I keratin, keratins 9 to 23) and a basic keratin (or type II keratin, keratins 1 to 8) assemble to form filaments. Keratin isoforms demonstrate tissue and differentiation-specific profiles that make them useful as research biomarkers. Research studies have shown that mutations in keratin genes are associated with skin disorders, liver and pancreatic diseases, and inflammatory intestinal

diseases.






Synonyms: Cell proliferation-inducing gene 46 protein; CK-18; CYK18; Cytokeratin-18; K18; Keratin, type I

cytoskeletal 18; Keratin-18; KRT18; PIG46

Note: For research use only, not for use in diagnostic procedure.

Product images:



p-Cytokeratin 8 (S52) p-Cytokeratin 18 (S52) Western blot (WB) analysis of p-Cytokeratin 18 (S52) polyclonal antibody at 1:500 dilution Lane1:HepG2 whole cell lysate(40µg) Lane2:L02 whole cell lysate(40µg) Lane3:A549 whole cell lysate(40µg)