

Product datasheet for TA392959S

OriGene Technologies, Inc.

9620 Medical Center Drive, Ste 200 Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com CN: techsupport@origene.cn

hnRNP K (HNRNPK) Rabbit Polyclonal Antibody

Product data:

Product Type: Primary Antibodies

Applications: WB

Reactivity: WB: 1:500~1:1000 Human, Mouse, Rat

Host: Rabbit Isotype: IgG

Clonality: Polyclonal

Immunogen: Synthetic phosphopeptide derived from human hnRNP K around the phosphorylation site of

Serine 284

Specificity: p-hnRNP K (S284) pAb detects endogenous levels of hnRNP K protein only when

phosphorylated at Ser284.

Formulation: Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

Predicted Protein Size: ~ 58 to 62 kDa

Gene Name: heterogeneous nuclear ribonucleoprotein K

Database Link: Entrez Gene 3190 Human

P61978

Background: The majority of hnRNP proteins components are localized to the nucleus; however some

shuttle between the nucleus and the cytoplasm, such as hnRNP K. hnRNP K, a 65 kDa protein, recruits a variety of molecular partners through two K homologous (KH) domains, which are

required for protein-protein interactions. hnRNP K also contains several potential

phosphorylation sites, including Ser 302, the major site of PKCd phosphorylation, which are thought to regulate various cellular functions, including sequence-specific DNA binding,

transcription, RNA binding and nucleocytoplasmic shuttling.






hnRNP K (HNRNPK) Rabbit Polyclonal Antibody - TA392959S

Synonyms: Heterogeneous nuclear ribonucleoprotein K; HNRNPK; hnRNP K; HNRPK; Transformation up-

regulated nuclear protein; TUNP

Note: For research use only, not for use in diagnostic procedure.

Product images:



p-hnRNP K (S284)

Western blot (WB) analysis of p-hnRNP K (S284) pAb at 1:500 dilution Lane1:Hela whole cell lysate(40µg) Lane2:HEPG2 whole cell lysate(40µg) Lane4:C6 whole cell lysate(40µg)