

## **Product datasheet for TA392965M**

## **CD11b (ITGAM) Mouse Polyclonal Antibody**

## **Product data:**

**Product Type:** Primary Antibodies

Applications: WE

**Reactivity:** WB: 1:500~1:1000 Human, Mouse, Rat

Host: Mouse Isotype: IgG

Clonality: Polyclonal

Immunogen: Recombinant Human CD11B (630aa-1104aa).

**Specificity:** CD11B (L1) mAb detects endogenous levels of CD11B protein. **Formulation:** 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Concentration: 1mg/ml

Conjugation: Unconjugated

Storage: Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.

Stability: 1 year

**Predicted Protein Size:** ~ 170 kDa

Gene Name: integrin subunit alpha M

Database Link: Entrez Gene 3684 Human

P11215

**Background:** Integrin αM, also designated complement component receptor-3 α, CD11b (p170),

macrophage antigen a polypeptide, cell surface glycoprotein Mac-1 a subunit, MAC1A, MO1A and ITGAM) is a cell adhesion molecule that acts as a receptor for cell surface ligands such as intracellular adhesion molecules (ICAMs) or soluble ligands. Integrins are heterodimeric proteins that contain an a chain and b chain. Integrin  $\alpha M$  combines with the Integrin  $\beta 2$  to form a leukocyte-specific integrin referred to as macrophage receptor 1 (Mac-1), or inactivated-C3b (iC3b) receptor 3 (CR3). Integrin  $\alpha M/\beta 2$  is important in the adherence of neutrophils and monocytes to stimulated endothelium, and also in the phagocytosis of

complement coated particles.



**OriGene Technologies, Inc.** 9620 Medical Center Drive, Ste 200

CN: techsupport@origene.cn

Rockville, MD 20850, US Phone: +1-888-267-4436 https://www.origene.com techsupport@origene.com EU: info-de@origene.com




Synonyms:

CD11 antigen-like family member B; CD11B; CD11b; Cell surface glycoprotein MAC-1 subunit alpha; CR-3 alpha chain; CR-3 $\alpha$ ; CR3 $\alpha$ ; CR3 $\alpha$ ; Integrin alpha-M; Integrin  $\alpha$ M; ITGAM; Leukocyte adhesion receptor MO1; Neutrophil adherence receptor

Note:

For research use only, not for use in diagnostic procedure.

## **Product images:**



Western blot (WB) analysis of CD11B (L1) mAb at 1:500 dilution Lane1:Myla2059 whole cell lysate(30ug) Lane2:HuT78 whole cell lysate(30ug) Lane3:K562 whole cell lysate(30ug) Lane4:The Spleen tissue lysate of Rat(40ug) Lane5:RAW264.7 whole cell lysate(40ug) Lane6:THP-1 whole cell lysate(40ug)