

Product datasheet for **TA396635**

GPD1 Goat Polyclonal Antibody

Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	WB: 1:500 - 1:3,000 ELISA: 1:5,000 - 1:25,000
Reactivity:	Rabbit
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Glycerol-3-Phosphate Dehydrogenase [Rabbit Muscle]
Specificity:	Anti-GLYCEROL-3-PHOSPHATE DEHYDROGENASE (GOAT) Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Glycerol-3-Phosphate Dehydrogenase [Rabbit Muscle]. Cross reactivity against Glycerol-3-Phosphate Dehydrogenase from other sources is unknown.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Reconstitution Method:	Restore with deionized water (or equivalent) - Reconstitution Volume: 100 µL
Concentration:	1.0 mg/ml - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at 4° C prior to restoration. For extended storage aliquot contents and freeze at -20° C or below. Avoid cycles of freezing and thawing. Centrifuge product if not completely clear after standing at room temperature. This product is stable for several weeks at 4° C as an undiluted liquid. Dilute only prior to immediate use.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	<u>Entrez Gene 100339469 Rabbit P08507</u>



[View online »](#)

Background:	Glycerol-3-phosphate dehydrogenase (GPDH) is an enzyme that catalyzes the reversible redox conversion of dihydroxyacetone phosphate (aka glycerone phosphate, outdated) to sn-glycerol 3-phosphate. Glycerol-3-phosphate dehydrogenase serves as a major link between carbohydrate metabolism and lipid metabolism. It is also a major contributor of electrons to the electron transport chain in the mitochondria. Older terms for glycerol-3-phosphate dehydrogenase include alpha glycerol-3-phosphate dehydrogenase (alphaGPDH) and glycerolphosphate dehydrogenase (GPDH). However, glycerol-3-phosphate dehydrogenase is not the same as glyceraldehyde 3-phosphate dehydrogenase (GAPDH) whose substrate is an aldehyde not an alcohol. Anti-Glycerol-3-Phosphate Dehydrogenase Antibody is ideal for investigators involved in glucose energy metabolism research.
Synonyms:	goat anti-Glycerol-3-Phosphate Dehydrogenase Antibody, FLJ26652 antibody, G3PD antibody, Gdc-1 antibody, Glycerphosphate dehydrogenase antibody, GPD-C antibody, Gpd1 protein antibody
Note:	Anti-GLYCEROL-3-PHOSPHATE DEHYDROGENASE (GOAT) Antibody has been assayed against 1.0 ug of Glycerol-3-Phosphate Dehydrogenase [Rabbit Muscle] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product. Specific conditions should be optimized by researcher.