

## Product datasheet for **TA396636S**

### GPD1 Goat Polyclonal Antibody

#### Product data:

Product Type:	Primary Antibodies
Applications:	ELISA, WB
Recommended Dilution:	<b>WB:</b> 1:500 - 1:3,000 <b>ELISA:</b> 1:5,000 - 1:25,000
Reactivity:	Rabbit
Host:	Goat
Clonality:	Polyclonal
Immunogen:	Glycerol-3-Phosphate Dehydrogenase [Rabbit Muscle]
Specificity:	Anti-GLYCEROL-3-PHOSPHATE DEHYDROGENASE (GOAT) Antibody is an IgG fraction antibody purified from monospecific antiserum by a multi-step process which includes delipidation, salt fractionation and ion exchange chromatography followed by extensive dialysis against the buffer stated above. Assay by immunoelectrophoresis resulted in a single precipitin arc against anti-Goat Serum as well as purified and partially purified Glycerol-3-Phosphate Dehydrogenase [Rabbit Muscle]. Cross reactivity against Glycerol-3-Phosphate Dehydrogenase from other sources is unknown.
Formulation:	0.02 M Potassium Phosphate, 0.15 M Sodium Chloride, pH 7.2
Concentration:	1.0 mg/ml - lot specific
Conjugation:	Unconjugated
Storage:	Store vial at -20° C or below prior to opening. This vial contains a relatively low volume of reagent (25 µL). To minimize loss of volume dilute 1:10 by adding 225 µL of the buffer stated above directly to the vial. Recap, mix thoroughly and briefly centrifuge to collect the volume at the bottom of the vial. Use this intermediate dilution when calculating final dilutions as recommended below. Store the vial at -20°C or below after dilution. Avoid cycles of freezing and thawing.
Stability:	Expiration date is one (1) year from date of receipt.
Database Link:	<a href="#">Entrez Gene 100339469 Rabbit P08507</a>



[View online »](#)

- Background:** Glycerol-3-phosphate dehydrogenase (GPDH) is an enzyme that catalyzes the reversible redox conversion of dihydroxyacetone phosphate (aka glycerone phosphate, outdated) to sn-glycerol 3-phosphate. Glycerol-3-phosphate dehydrogenase serves as a major link between carbohydrate metabolism and lipid metabolism. It is also a major contributor of electrons to the electron transport chain in the mitochondria. Older terms for glycerol-3-phosphate dehydrogenase include alpha glycerol-3-phosphate dehydrogenase (alphaGPDH) and glycerolphosphate dehydrogenase (GPDH). However, glycerol-3-phosphate dehydrogenase is not the same as glyceraldehyde 3-phosphate dehydrogenase (GAPDH) whose substrate is an aldehyde not an alcohol. Anti-Glycerol-3-Phosphate Dehydrogenase Antibody is ideal for investigators involved in glucose energy metabolism research.
- Synonyms:** goat anti-Glycerol-3-Phosphate Dehydrogenase Antibody, FLJ26652 antibody, G3PD antibody, Gdc-1 antibody, Glycerphosphate dehydrogenase antibody, GPD-C antibody, Gpd1 protein antibody
- Note:** Anti-GLYCEROL-3-PHOSPHATE DEHYDROGENASE (GOAT) Antibody has been assayed against 1.0 ug of Glycerol-3-Phosphate Dehydrogenase [Rabbit Muscle] in a standard ELISA using Peroxidase conjugated Affinity Purified anti-Goat IgG [H&L] (Goat) code #611-1302 and (ABTS (2,2'-azino-bis-[3-ethylbenthiazoline-6-sulfonic acid]) code # ABTS-100 as a substrate for 30 minutes at room temperature. A working dilution of 1:20,000 to 1:100,000 of the reconstitution concentration is suggested for this product. Specific conditions should be optimized by researcher.